

AD-A045 864

ILLINOIS UNIV AT URBANA-CHAMPAIGN DEPT OF CHEMISTRY

F/G 7/3

DIOXETANONE CHEMILUMINESCENCE BY THE CHEMICALLY INITIATED ELECT--ETC(U)

OCT 77 S P SCHMIDT, G B SCHUSTER

N00014-76-C-0745-07

NL

UNCLASSIFIED

1 of 1
ADA045864



Microfiche frame containing text

Microfiche frame containing text

END
DATE
FILMED
11-77
DDC

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE

READ INSTRUCTIONS
BEFORE COMPLETING FORM

1. REPORT NUMBER N0014-76-C-0745-07	2. GOVT ACCESSION 11	3. RECIPIENT'S CATALOG NUMBER 11
4. TITLE (and Subtitle) Dioxetanone Chemiluminescence by the Chemically Initiated Electron Exchange Pathway. The Efficiency Generation of Excited Singlet States.	5. TYPE OF REPORT & PERIOD COVERED Technical	
7. AUTHOR(s) Gary B. Schuster Steven P. Schmidt	6. PERFORMING ORG. REPORT NUMBER	8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0745
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Illinois Urbana, Illinois 61801	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR-051-616	
11. CONTROLLING OFFICE NAME AND ADDRESS Chemistry Program, Materials Science Division, Office of Naval Research, 800 N. Quincy St., Arlington, VA 22217	12. REPORT DATE October 14, 1977	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. NUMBER OF PAGES 11	
	15. SECURITY CLASS. (of this Report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for Public Release, Distribution Unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) DDC RECEIVED OCT 26 1977 F.		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) chemiluminescence dioxetane electron transfer firefly		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The chemiluminescence of dimethyldioxetanone (1) in the presence of easily oxidized aromatic hydrocarbons was studied. It was observed that the rate of reaction of (1) depends upon the nature and concentration of the aromatic hydrocarbon. The efficiency of light production was correlated with the one electron oxidation potential of the hydrocarbon. These results are interpreted in terms of the recently described chemically initiated electron exchange luminescence. drawing on back		

DD FORM 1473


EDITION OF 1 NOV 63 IS OBSOLETE
S/N 0102-14-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

AD A 045864

DDC FILE COPY

20. Abstract



OFFICE OF NAVAL RESEARCH

Contract ^{N00014-} N0014-76-C-0745

Task No. NR-051-616

TECHNICAL REPORT NO. ^{N00014-} N0014-76-C-0745-07

6 Dioxetanone Chemiluminescence by the Chemically Initiated Electron
Exchange Pathway. The Efficient Generation of Excited Singlet States.

by

10 Steven P. Schmidt and Gary B. Schuster

Prepared for Publication

in

Journal of the American Chemical Society

School of Chemical Sciences
University of Illinois
Urbana, Illinois 61801

11 14 October 1977

12 13 p.


Reproduction in whole or in part is permitted for
any purpose of the United States Government

Approved for Public Release; Distribution Unlimited.

408 087

Dioxetanone Chemiluminescence by the Chemically Initiated Electron Exchange Pathway. The Efficient Generation of Excited Singlet States.¹

Abstract: The chemiluminescence of dimethyldioxetanone (1) in the presence of easily oxidized aromatic hydrocarbons was studied. It was observed that the rate of reaction of 1 depends upon the nature and concentration of the aromatic hydrocarbon. The efficiency of light production was correlated with the one electron oxidation potential of the hydrocarbon. These results are interpreted in terms of the recently described chemically initiated electron exchange luminescence.




ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
	SPECIAL
<div style="font-size: 2em; font-weight: bold; text-align: center;">A</div>	

Sir:

Many recent studies of chemiluminescence from organic molecules have centered on the reactions of the strained four membered ring peroxides known as dioxetanes.² These molecules have been identified or implicated in many of the most efficient chemi- and bioluminescent reactions. In particular, a carbonyl substituted dioxetane (dioxetanone) has been identified as the key high energy molecule responsible for light production in the firefly.³ In this paper we report our findings on the mechanism of light production from simply substituted dioxetanones. Our studies show that the most important light forming path for reaction of dioxetanone **1** is the bimolecular route we have recently identified as chemically initiated electron exchange luminescence (CIEEL).⁴ Moreover, we have found that under conditions favoring CIEEL the fraction of reacting dioxetanone molecules that generate a photon of light approaches the most efficient bioluminescent reactions known. In addition, by analogy, it appears that the initiating reaction in firefly bioluminescence is an intramolecular electron transfer akin to the observed intermolecular reaction reported herein.⁵

Dimethyldioxetanone (**1**) was prepared and purified according to the procedure of Adam.⁶ Thermolysis of **1** in CH_2Cl_2 at 24.5° leads to the quantitative generation of acetone and to light emission.⁷ The observed chemiluminescence under these conditions is a result of acetone emission. It has been reported previously that the addition of certain aromatic hydrocarbons to solutions of dioxetanone **1** results in the generation of hydrocarbon luminescence and markedly increases the light yield.⁸ We have also observed this effect and, significantly, have found that the aromatic hydrocarbon also increases the rate of reaction of the dioxetanone.

The rate acceleration is directly proportional to the concentration of the hydrocarbon (catalytic chemiluminescence activator), as is shown for rubrene in Figure 1.



Also, the observed rate acceleration is dependent on the structure of the activator. These findings are inconsistent with the previously proposed unimolecular pathway for excited state production. In the original mechanism, thermal reaction of the dioxetanone generated electronically excited acetone which, in a subsequent step, transferred energy to the hydrocarbon. If this scheme was operative, the rate of reaction of the dioxetanone would be independent of the structure and concentration of the added hydrocarbon. This is contrary to our results.

To probe the nature of the interaction between dimethyldioxetanone (1) and the catalytic chemiluminescence activator, the efficiency of generation of light with various activators was examined. We observed that at identical hydrocarbon concentrations there is nearly a hundred fold range in the ability of the different activators to catalyze excited state formation.⁹ This finding indicates that it is the bimolecular reaction between the dioxetanone and the activator that is responsible for the major fraction of the hydrocarbon chemiluminescence. Critically, the only predictor of activator efficiency is the one electron oxidation potential of the hydrocarbon, as is shown in Figure 2. This result is required if electron transfer from the activator to the dioxetanone is the rate determining step for formation of the activator excited singlet state. These findings are entirely consistent

with excited state production from dioxetanone I^* by the mechanism we have recently identified as chemically initiated electron exchange luminescence.⁴

The efficiency of production of excited singlet activator by the induced decomposition of I^* was investigated by measuring the absolute chemiluminescence yield.¹⁰ When rubrene was employed as the activator in CH_2Cl_2 solution at 24.5° it was found that $17 \pm 5\%$ of the dioxetanone molecules that proceed through the bimolecular path generate an excited rubrene singlet state. Even though this system is unoptimized, the remarkably high efficiency approaches that of the known bioluminescent reactions.¹²


The proposed mechanism for formation of the emitting singlet state of the chemiluminescence activator by the CIEEL path is shown in Scheme 1. The first step is a one electron transfer from the activator to the dioxetanone. The rate of this process, of course, depends upon the activation barrier for the reaction which is determined, in part, by the oxidation potential of the activator. Similiar electron transfer reactions have been postulated, for example, to account for the catalytic induced decomposition of peroxides by transition metals and amines.¹³ The activation energy for the electron transfer is also dependent upon the reduction potential of the peroxide. For dioxetanone I^* the bimolecular rate constant for the CIEEL path (k_2) in CH_2Cl_2 at 24.5° with rubrene is $0.44 \text{ M}^{-1} \text{ s}^{-1}$. This compares to a value of $8 \text{ M}^{-1} \text{ s}^{-1}$ determined for diphenoyl peroxide under similar conditions.⁴ The difference in the catalytic rate constants for these two peroxides is predicted by their reduction potentials.¹⁴ The electron withdrawing carbonyl groups flanking the oxygen-oxygen bond of the diacyl peroxide results in more facile reduction and concomitantly a larger k_2 for diphenoyl peroxide.

The next step along the chemiluminescence path is the rapid loss of CO_2 from the reduced dioxetanone. This generates the radical anion of acetone within the same solvent cage as the radical cation of the activator.¹⁵

Subsequent charge annihilation of the cage radical ions results in electronically excited state generation.¹⁶ One explanation for the observed 17% singlet yield is that the decarboxylation and charge annihilation are competitive with spin equilibration. Thus, the dioxetanones that react by the CIEEL path, but do not generate an excited singlet activator, may be generating the undetected triplet excited state of the hydrocarbon. The final step in the sequence is light emission from the excited activator, which we detect as chemiluminescence.

In competition with the CIEEL path, uncatalyzed unimolecular decomposition of the dioxetanone generates electronically excited acetone. The combination of these two excitation mechanisms accounts for all of the experimental observations on the chemiluminescence of dioxetanone **1**.

Scheme I



In summary, we have shown that an efficient CIEEL pathway is the major light generating process from dioxetanone **1** with any one of several easily oxidized activators. This is the third example of efficient chemiluminescence by this route.^{4,17} We are continuing our investigation of the chemiluminescence of dioxetanones to further establish the details of the mechanism in this case. We are also investigating other chemiluminescent systems that appear to react by the CIEEL path.

Acknowledgment

This work was supported in part by the Office of Naval Research and in part by the donors of the Petroleum Research Fund administered by the American Chemical Society. We wish to thank Mr. William Wokas for the preparation of 9,10-diphenylethynylanthracene.

Steven P. Schmidt and Gary B. Schuster*¹⁸

Department of Chemistry
Roger Adams Laboratory
University of Illinois
Urbana, Illinois 61801

References


1. These results were presented in part at the 1977 annual meeting of the American Chemical Society, Chicago, Illinois, August 28, 1977.
2. For leading references see: T. Wilson, Int. Rev. Sci.: Phys. Chem., Ser. Two, 9, 265 (1976).
3. E. H. White, J. D. Miano, and M. Umbreit, J. Am. Chem. Soc., 97, 198 (1975).
4. J-y. Koo and G. B. Schuster, J. Am. Chem. Soc., 99, 6107 (1977).
5. This point has been discussed more fully, J-y. Koo, S. P. Schmidt, and G. B. Schuster, Proc. Natl. Acad. Sci. U. S., in press.
6. W. Adam and O. Cueto, J. Org. Chem., 42, 38 (1977); W. Adam and J.-C. Liu, J. Am. Chem. Soc., 94, 2894 (1972); W. Adam, A. Alzerreca, J.-C. Liu, and F. Yany, J. Am. Chem. Soc., 99, 5768 (1977).
7. Typical conditions; $[I] = 10^{-4}$ M in CH_2Cl_2 at 24.5° . Light was detected by the photon counting technique. Spectral resolution was achieved with a Jarrel-Ash 0.25 M monochromator.
8. W. Adam, G. A. Simpson, and F. Yany, J. Phys. Chem., 78, 2559 (1974).
9. The intensity was determined at 2×10^{-5} M in activator in CH_2Cl_2 solution at 24.5° under low resolution conditions so that the emission spectra were Gaussian. The intensities were corrected for photomultiplier tube and monochromator efficiencies according to data supplied by the manufacturers. All solutions contained 20 μ l of 5% aqueous Na_4EDTA to suppress metal catalyzed reactions.
10. The yield of light was determined relative to tetramethyldioxetane (TMD) using 9,10-dibromoanthracene as the acceptor. The yield of acetone triplet was taken to be 30%,^{11a} the triplet-singlet energy transfer efficiency at 40%,^{11b} and the fluorescence quantum yield to be 10%.


11. a) T. Wilson, D. E. Golan, M. S. Harris, and A. L. Baumstark, J. Am. Chem. Soc., 98, 1086 (1976); b) R. Schmidt, H. Kelm, and H.-D. Brauer, Ber. Bunsenges Phys. Chem., 81, 402 (1977).
12. Firefly bioluminescence is claimed to be 80% efficient; W. D. McElroy, Fedn. Proc. Fedn. Am. Soc. Exp. Biol., 19, 941 (1960).
13. For example see: J. K. Kochi and H. E. Mains, J. Org. Chem., 30, 1862 (1965).
14. Peroxyesters typically have reduction potentials of ca. -0.9 eV while diacyl peroxides reduce at ca. -0.1 eV versus the saturated calomel electrode: L. S. Silbert, L. P. Witnauer, D. Swern, and C. Ricciati, J. Am. Chem. Soc., 81, 3244 (1959).
15. Air saturated solutions were nearly as efficient as argon purged solutions. Free radical anions should react rapidly with O₂.
16. L. R. Faulkner, Int. Rev. Sci.: Phys. Chem., Ser. Two, 9, 213 (1977).
17. J. P. Smith and G. B. Schuster, J. Am. Chem. Soc., submitted.
18. Fellow of the Alfred P. Sloan Foundation.

Captions for Figures

Figure 1. Observed dependence of rate of reaction for dioxetanone on the rubrene concentration. Note that the extrapolated zero rubrene concentration rate agrees, within experimental error, to the independently determined value.

Figure 2. Dependence of the chemiluminescence intensity by the CIEEL path on oxidation potential (E_{ox}) of the activator. In order of increasing oxidation potential the points are: rubrene, perylene, 9,10-diphenylethynylanthracene, and 9,10-diphenylanthracene.





ONR TECHNICAL REPORT DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research ²		Commander	
Arlington, Virginia 22217		Naval Undersea Research & Development	
Attn: Code 472	2	Center	
Office of Naval Research		San Diego, California 92132	
Arlington, Virginia 22217		Attn: Technical Library, Code 133	1
Attn: Code 102IP	6	Naval Weapons Center	
ONR Branch Office		China Lake, California 93555	
536 S. Clark Street		Attn: Head, Chemistry Division	1
Chicago, Illinois 60605		Naval Civil Engineering Laboratory	
Attn: Dr. George Sandoz	1	Port Hueneme, California 93041	
ONR Branch Office		Attn: Mr. W. S. Haynes	1
715 Broadway		Professor O. Heinz	
New York, New York 10003		Department of Physics & Chemistry	
Attn: Scientific Dept.	1	Naval Postgraduate School	
ONR Branch Office		Monterey, California 93940	1
1030 East Green Street		Dr. A. L. Slafkosky	
Pasadena, California 91106		Scientific Advisor	
Attn: Dr. R. J. Marcus	1	Commandant of the Marine Corps (Code RD-1)	
ONR Branch Office		Washington, D.C. 20380	1
760 Market Street, Rm. 447		Dr. M. A. El-Sayed	
San Francisco, California 94102		University of California	
Attn: Dr. P. A. Miller	1	Department of Chemistry	
ONR Branch Office		Los Angeles, California 90024	1
495 Summer Street		Dr. M. W. Windsor	
Boston, Massachusetts 02210		Washington State University	
Attn: Dr. L. H. Peebles	1	Department of Chemistry	
Director, Naval Research Laboratory		Pullman, Washington 99163	1
Washington, D.C. 20390		Dr. E. R. Bernstein	
Attn: Library, Code 2029 (ONRL)	6	Colorado State University	
Technical Info. Div.	1	Department of Chemistry	
Code 6100, 6170	1	Fort Collins, Colorado 80521	1
The Asst. Secretary of the Navy (R&D)		Dr. C. A. Heller	
Department of the Navy		Naval Weapons Center	
Room 4E736, Pentagon		Code 6059	
Washington, D.C. 20350	1	China Lake, California 93555	1
Commander, Naval Air Systems Command		Dr. G. Jones, II	
Department of the Navy		Boston University	
Washington, D.C. 20360		Department of Chemistry	
Attn: Code 310C (H. Rosenwasser)	1	Boston, Massachusetts 02215	1
Defense Documentation Center		Dr. M. H. Chisholm	
Building 5, Cameron Station		Chemistry Department	
Alexandria, Virginia 22314	12	Princeton, New Jersey 08540	1
U.S. Army Research Office		Dr. J. R. MacDonald	
P.O. Box 12211		Code 6110	
Research Triangle Park, North Carolina		Chemistry Division	
Attn: CRD-AA-IP	1	Naval Research Laboratory	
		Washington, D.C. 20375	1

ONR TECHNICAL REPORT DISTRIBUTION LIST

No. Copies

Dr. G. B. Schuster Chemistry Department University of Illinois Urbana, Illinois 61801	1
Dr. E. M. Eyring University of Utah Department of Chemistry Salt Lake City, Utah	1
Dr. A. Adamson University of Southern California Department of Chemistry Los Angeles, California 90007	1
Dr. M. S. Wrighton Massachusetts Institute of Technology Department of Chemistry Cambridge, Massachusetts 02139	1